



ELSEVIER

Journal of Pure and Applied Algebra 177 (2003) 1–4

JOURNAL OF
PURE AND
APPLIED ALGEBRAwww.elsevier.com/locate/jpaa

Writing certain commutators as products of cubes in free groups [☆]

Mehri Akhavan-Malayeri^{a,b}^a*Department of Mathematics, Azzahra University, Vanak, Tehran, 19834, Iran*^b*Institute for Studies in Theoretical Physics and Mathematics, Niavaran Square, P.O. Box 19395-1795, Tehran, Iran*

Received 10 February 2002

Communicated by C.A. Weibel

Abstract

Let $\text{Cu}(\gamma)$ be the minimal number of cubes required to express an element γ of a free group F . We establish a method for showing that certain equations do not have solutions in free groups. Using it, we find $\text{Cu}(\gamma)$ for certain elements of the derived subgroup of F . If $W = F \wr C_\infty$ is the wreath product of F by the infinite cyclic group, we also show that every element of W' is a product of at most one commutator and three cubes in W . © 2002 Elsevier Science B.V. All rights reserved.

MSC: 20F12; 20F99

1. Introduction

It is known that conjugate elements commute in a group of exponent 3. In [2], Lyndon wrote $[x^y, y]$ as a product of four cubes:

$$[x^y, x] = (y^{-1}x^{-1}y^2)^3 y^{-3} (yx)^3 (x^{-1}y^{-1}x^{-1}y^{-1}xyx)^3.$$

This is not optimal. We show that $[x^y, x]$ can be expressed as a product of 3 cubes, but not fewer than 3. We use the following notation:

If γ is an element of a group G , we write $\gamma \in \text{Cu}_n = \text{Cu}_n(G)$ to mean that γ can be written as a product of n cubes in G , and $\text{Cu}(\gamma) = n$ denotes that $\gamma \in \text{Cu}_n$ but $\gamma \notin \text{Cu}_{n-1}$. We call $\text{Cu}(\gamma)$ the cube length of γ . Finally, G^3 is the fully invariant subgroup of G , generated by the cubes of the elements of G .

[☆] This research was in part supported by a grant from IPM.

E-mail address: mmalayer@azzahra.ac.ir, makhavanm@yahoo.com (M. Akhavan-Malayeri).

2. Main results

Let F be a free group and x, y be two distinct elements of a free generating set. Then $[x^y, x] = [x, y, x]$. We present the following easily verified expression of $[x, y, x]$ as a product of three cubes;

$$[x, y, x] = ((x^{-1}y)^y)^3 (y^{-2}x)(y^x)^3. \quad (1)$$

Hence, $\text{Cu}([x, y, x]) \leq 3$. To show that 3 is the minimal number, we consider equation $[x, y, x] = A^3 B^3$ in F . We reduce the problem to the rank two case, we may assume $F = F(x, y)$ be the free group of rank 2 generated by x, y , and we prove the following results:

Theorem 1. Let $F(x, y)$ be the free group of rank 2 freely generated by x, y , the equation $[x, y, x]^n = A^3 B^3$ has a solution in F if and only if $3|n$. In particular, neither $[x, y, x] = A^3 B^3$ nor $[x, y, x]^2 = A^3 B^3$ has a solution.

To prove this, we need the following lemma. Let F_n denote the n th term of the lower central series of F .

Lemma 1. Let $F = F(x, y)$ be a free group, and $\gamma \in \text{Cu}_2(F)$. If $\gamma \in F_n$ for some n then the image of γ in F_n/F_{n+1} belongs to $\text{Cu}_1(F_n/F_{n+1})$ and the image of γ in F_n/F_{n+2} belongs to $\text{Cu}_2(F_n/F_{n+1})$.

We prove following theorem regarding the wreath product of groups.

Theorem 2. Let $W = F \wr C_\infty$ be the wreath product of a free group F and the infinite cyclic group C_∞ . Then every element of W' is a product of at most one commutator and three cubes in W . More precisely let B be the base group of W and let t generate C_∞ . For any $w \in W'$ there is $a \in B$ and $c_1, c_2, c_3 \in W$ such that

$$w = [a, t]c_1^3 c_2^3 c_3^3.$$

3. Proofs

Proof of Lemma 1. If $n = 1$, $\gamma \in F_1$ hence $\gamma = u^3 v^3$ in F . Since $\gamma F' = u^3 v^3 F' = (uv)^3 F'$, $\gamma = (uv)^3$ modulo F' . When $n > 1$, $\gamma \in F'$, and $(uv)^3 = 1 \pmod{F'}$. Since F/F' is torsion free $uv = f \in F'$. Hence $\gamma = u^3 v^3 = u^3 (u^{-1}f)^3 = u^2 f u^{-1} f u^{-1} f = [u^{-1}, f^{-1}]^{u^{-1}} [u^{-1}, f^{-2}] f^3$. But then $\gamma \equiv f^3 \pmod{F_3}$.

Also, we have

$$\begin{aligned} \gamma &\equiv [u^{-1}, f^{-1}] [u^{-1}, f^{-1}, u^{-1}] [u^{-1}, f^{-1}]^2 [u^{-1}, f^{-1}, f^{-1}] f^3 \\ &\equiv [u^{-1}, f^{-1}]^3 f^3 \pmod{F_4}. \end{aligned}$$

Therefore, if $\gamma \in F' = F_2$ and $\gamma \in \text{Cu}_2(F)$ then the image of γ in F_2/F_3 belongs to $\text{Cu}_1(F_2/F_3)$ and the image of γ in F_2/F_4 belongs to $\text{Cu}_2(F_2/F_4)$.

We show if $\gamma \in F_n$ then $f = uv \in F_n$. Let $\gamma = u^3v^3 \in F_n$ and suppose $uv \notin F_n$. Assume $r \leq n$ be the least integer such that $uv \in F_{r-1} - F_r$. Then $(uv)^3 \in F_{r-1} - F_r$, because F_{r-1}/F_r is torsion free. Hence, $u^3v^3 \in F_{r-1} - F_r$ and $\gamma \notin F_r$, a contradiction.

Suppose now that $\gamma \in F_n$ ($n \geq 2$). Then $f = uv \in F_n$, hence $[u^{-1}, f^{-1}]$ and $[u^{-1}, f^{-2}]$ belong to F_{n+1} and

$$\gamma = [u^{-1}, f^{-1}]^{u^{-1}} [u^{-1}, f^{-2}] f^3 \equiv f^3 \pmod{F_{n+1}}.$$

From the last equation we have

$$\begin{aligned} \gamma &= [u^{-1}, f^{-1}] [u^{-1}, f^{-1}, u^{-1}] [u^{-1}, f^{-1}]^2 [u^{-1}, f^{-1}, f^{-1}] f^3 \\ &\equiv [u^{-1}, f^{-1}]^3 f^3 \pmod{F_{n+2}} \end{aligned}$$

and the proof is complete. Now we turn to the proof of Theorem 1. \square

Proof of Theorem 1. Let $[x, y, x]^n = A^3B^3$ in F . By Lemma 1 there is a $u \in F_3$, such that $[x, y, x]^n \equiv u^3 \pmod{F_4}$. Since $[x, y, x]$ belongs to a basis of the free group F_2/F_3 , it follows that $[x, y, x]^n \equiv u^3 \pmod{F_4}$ iff $3|n$. This completes the proof of Theorem 1. \square

Corollary 1. Let $\gamma = [c_{i_1}, \dots, c_{i_k}, c_{i_{k+1}}, c_{i_{k+1}}]$ be a basic commutator in F . Then $\text{Cu}(\gamma) = 3$, in particular if $\gamma = [x, y, x]$ then $\text{Cu}(\gamma) = 3$.

Proof. Set $\omega = [c_{i_1}, \dots, c_{i_k}]$, so that $\gamma = [[\omega, c_{i_{k+1}}], c_{i_{k+1}}]$. Then by Eq. (1), $\gamma \in \text{Cu}_3(F)$. By Lemma 1, $\gamma^n = A^3B^3$ implies that $\gamma^n \equiv u^3 \pmod{F_{k+3}}$, where u is an element of F_{k+2} . Since γ is a basic commutator, it follows that $\gamma^n \equiv u^3 \pmod{F_{k+3}}$ iff $3|n$, hence $\text{Cu}(\gamma) \neq 2$. In particular, by Theorem 1 and Eq. (1), $\text{Cu}[x, y, x] = 3$. \square

Corollary 2. With the hypothesis of Corollary 1, $3 \leq \text{Cu}(\gamma^2) \leq 4$, in particular $3 \leq \text{Cu}[x, y, x]^2 \leq 4$.

Proof. Since $\gamma^2 = \gamma^{-1}\gamma^3$, Corollary 1 implies that $\text{Cu}(\gamma^2) \leq 4$. By Theorem 1 and the proof of Corollary 1, $3 \leq \text{Cu}(\gamma^2)$. The proof is complete. \square

Proof of Theorem 2. In [1] we proved that every element of W' is a product of at most three commutators in W . Let B be the base group of W and let $T = \langle t \rangle \cong C_\infty$. Then $W = BT$ is the semidirect product of B and T and $W' = [B, T] = \{[a, t], a \in B\}$. Similarly, $F_3(W) = [B, T, T] = \{[a, t, t], a \in B\}$, hence $\text{Cu}(F_3(W)) \leq 3$. According to [1] every element of W' modulo B' is a commutator, and if g is an arbitrary element of B' there are suitable elements say b, c in B such that

$$g = [b^{-1}, t^m][c^{-1}, t^{2m}] = (t^{-m})^{b-1} (t^{-2m})^{c-1} t^m t^{3m}.$$

Without loss of generality, we may assume m is a multiple of 3. Hence, $\text{Cu}(B') \leq 3$ and every element of W' can be written as a product of a commutator and three cubes in W . \square

Acknowledgements

I thank Professor A. Rhemtulla of the University of Alberta for his helpful suggestions regarding the preparation of this paper. I would like to thank the managing editor of the Journal of Pure and Applied Algebra and the referee who have patiently read and verified this note, and also suggested valuable comments. The author also like to acknowledge the supports of the institute for studies in Theoretical Physics and Mathematics and Azzahra University.

References

- [1] M. Akhavan-Malayeri, Commutator length and square length of the wreath product of a free group by the infinite cyclic group, *Houston J. Math.* 27 (4) (2001) 753–756.
- [2] R.C. Lyndon, Equations in groups, *Bol. Soc. Brasil. Mat.* 11 (1980) 79–102.